

Bilarge leptonic mixing from Abelian horizontal symmetries

Tommy Ohlsson^{a*}, Gerhart Seidl^{a†}

^a*Institut für Theoretische Physik, Physik-Department, Technische Universität München,
James-Frank-Straße, 85748 Garching bei München, Germany*

Abstract

We construct and present a model for leptonic mixing based on higher-dimensional operators, using the Froggatt–Nielsen mechanism, and Abelian horizontal symmetries (flavor symmetries) of continuous and discrete type. Our model naturally yields bilarge leptonic mixing, coming from both the charged leptons and the neutrinos, and an inverted neutrino mass hierarchy spectrum. The obtained values of the parameters, *i.e.*, the leptonic mixing parameters and the neutrino mass squared differences, are all consistent with the atmospheric neutrino data and the Mikheyev–Smirnov–Wolfenstein large mixing angle solution for the solar neutrino problem.

PACS: 14.60.Pq, 11.30.Hv

Key words: Leptonic mixing; Neutrino masses; Discrete symmetries; Higher-dimensional operators

1. Introduction

Predicting the pattern of fermion masses and mixings from a fundamental gauge theory is one of the major challenges in particle physics. In such an approach, the observed hierarchy of the fermion masses is usually understood in terms of some symmetry breaking interaction. In fact, due to their plausible Majorana nature, the extreme smallness of the neutrino masses could be associated with a violation of the $B - L$ symmetry. Thus, the neutrinos can shed light on the origin of the fermion masses and mixings, since most grand unified theories (GUTs) based on $SO(10)$ or E_6 and also string theories indeed expect the $B - L$ symmetry to be broken [1]. In building models, it is therefore particularly important to naturally reproduce the neutrino mass squared differences and the leptonic mixing parameters that have been determined by the atmospheric [2,3] and solar [4] neutrino data.

The neutrino mass squared differences are generally defined as

$$\Delta m_{ab}^2 \equiv m_a^2 - m_b^2,$$

where m_a is the mass of the a th neutrino mass eigenstate. We will here assume that there are three neutrino flavors, and therefore, three neutrino flavor states ν_α ($\alpha = e, \mu, \tau$) and also three

neutrino mass eigenstates ν_a ($a = 1, 2, 3$). The unitary leptonic mixing matrix¹ is then given by

$$U = (U_{\alpha a}) \equiv U^{\ell\dagger} U^\nu,$$

where the unitary mixing matrix U^ℓ (U^ν) rotates the left-handed charged lepton fields (the neutrino fields) so that the charged lepton mass matrix (the neutrino mass matrix) becomes diagonal. Thus, the leptonic mixing matrix acquires contributions from both the charged leptons and the neutrinos. These contributions usually add in a non-trivial way (see App. A). For three neutrino flavors, in the so-called standard parameterization, the leptonic mixing matrix reads

$$U = \begin{pmatrix} C_2 C_3 & S_3 C_2 & S_2 e^{-i\delta} \\ -S_3 C_1 - S_1 S_2 C_3 e^{i\delta} & C_1 C_3 - S_1 S_2 S_3 e^{i\delta} & S_1 C_2 \\ S_1 S_3 - S_2 C_1 C_3 e^{i\delta} & -S_1 C_3 - S_2 S_3 C_1 e^{i\delta} & C_1 C_2 \end{pmatrix},$$

where $S_a \equiv \sin \theta_a$, $C_a \equiv \cos \theta_a$ (for $a = 1, 2, 3$), and δ is the physical $\mathcal{CP}$ phase. Here $\theta_1 \equiv \theta_{23}$, $\theta_2 \equiv \theta_{13}$, and $\theta_3 \equiv \theta_{12}$ are the leptonic mixing angles. Recent results suggest that among the possible solutions to the solar neutrino problem, the Mikheyev–Smirnov–Wolfenstein (MSW) [6] large mixing angle (LMA) solution is somewhat preferred to the MSW small mixing angle (SMA) solution, the MSW low mass (LOW) solution, and the vacuum oscillation (VAC) solution [7–9]. Actually, a global solar two flavor neutrino oscillation analysis including the latest SNO

*E-mail: tohlsson@ph.tum.de

†E-mail: gseidl@ph.tum.de

¹The leptonic mixing matrix is sometimes called the Maki–Nakagawa–Sakata (MNS) mixing matrix [5].

data strongly favors the MSW LMA solution [10]. However, the MSW LMA solution excludes maximal solar mixing at the 95% confidence level [8] (and now even at the 99.73% confidence level [9]), and therefore, it also disfavors the scenario of so-called bimaximal mixing [11]. Thus, we will instead, most probably, have a *bilarge* mixing scenario in which the solar mixing angle θ_{12} is large, but not maximal, and the atmospheric mixing angle θ_{23} is approximately maximal. It is interesting to observe that there are very strong indications that the leptonic mixing is large, whereas, on the other hand, it has turned out experimentally that the quark mixing is small [12].

In this paper, we will investigate a model, which yields in a technically natural way bilarge leptonic mixing, reproduces the observed mass hierarchy of charged leptons, and leads to an inverted neutrino mass hierarchy spectrum. This will be achieved by generating lepton mass matrix textures, where the mixing of the charged leptons is comparable with the mixing of the quarks and the mixing of the neutrinos is essentially bimaximal. The striking difference between the bilarge leptonic mixing and the small quark mixing will then be accounted for by the neutrinos (mainly) and the charged leptons (partly).

This paper is organized as follows: In Sec. 2, we will introduce a model by adding to the standard model (SM) a set of extra fields and horizontal symmetries that will give rise to specific effective Yukawa interactions for the charged leptons. Next, minimizing the corresponding scalar potential, we will naturally obtain a mass matrix texture for the charged leptons, which is in agreement with experimental data. In Sec. 3, we will extend the representation content of the model in order to also obtain a realistic mass matrix texture for the neutrinos. (In App. A, we will explicitly calculate the leptonic mixing angles coming from the diagonalizations of the charged lepton and neutrino mass matrices, respectively.) Finally, in Sec. 4, we will present a summary as well as our conclusions.

2. Charged leptons

2.1. Horizontal symmetries

We will here consider an extension of the SM in which the lepton masses arise from higher-dimensional operators [13] via the Froggatt–Nielsen mechanism [14]. (For recent studies, see, *e.g.*, Ref. [15].) We will write, in a self-explanatory notation, the lepton doublets as $L_\alpha = (\nu_{\alpha L} \ e_{\alpha L})$, where $\alpha = e, \mu, \tau$, and the right-handed charged leptons as $E_\alpha = e_{\alpha R}$, where $\alpha = e, \mu, \tau$. Suppose that the part of the scalar sector, which transforms non-trivially under the SM gauge group, consists of two Higgs doublets H_1 and H_2 , where H_1 couples to the neutrinos and H_2 to the charged leptons.² (For simplicity and without loss of generality, the quark sector will be left out in our entire discussion.) Let us first restrict our discussion to the generation of the charged lepton masses. In order to obtain the structure of the charged lepton mass matrix from an underlying symmetry principle, we will further extend the scalar sector by SM singlet scalar fields ϕ_i ($i = 1, 2, \dots, 8$) and θ and assign the fields gauged horizontal U(1) charges Q_1, Q_2 , and Q_3 as follows:

	(Q_1, Q_2, Q_3)
L_e, E_e	$(1, 0, 0)$
$L_\mu, L_\tau, E_\mu, E_\tau$	$(0, 1, 0)$
ϕ_1, ϕ_2	$(-1, 1, 2)$
ϕ_3, ϕ_4	$(1, -1, 2)$
ϕ_5, ϕ_6	$(0, 0, 0)$
ϕ_7, ϕ_8	$(0, 0, 1)$
θ	$(0, 0, -1)$

In the rest of the paper, it is always understood that the Higgs doublets H_1 and H_2 are total singlets under transformations of the additional symmetries. Note that our model is kept anomaly-free, since the fermions transform as vector-like pairs under the extra U(1) charges. Next, the charges (Q_1, Q_2) of the charged lepton-

²This can easily be achieved by imposing a discrete $\mathbb{Z}_2$ symmetry under which H_2 and E_α ($\alpha = e, \mu, \tau$) are odd and H_1 and the rest of the SM fields are even.

antilepton pairs are given by:

		E_e	E_μ	E_τ
	(Q_1, Q_2)	$(1, 0)$	$(0, 1)$	$(0, 1)$
$\overline{L_e}$	$(-1, 0)$	$(0, 0)$	$(-1, 1)$	$(-1, 1)$
$\overline{L_\mu}$	$(0, -1)$	$(1, -1)$	$(0, 0)$	$(0, 0)$
$\overline{L_\tau}$	$(0, -1)$	$(1, -1)$	$(0, 0)$	$(0, 0)$

We observe that these charges forbid dimension-four Yukawa coupling terms for the coupling of the first generation to the second and third generations. A realistic charged lepton mass matrix will arise if we, in addition to the U(1) charges, introduce a set of discrete symmetries $\mathcal{D}_i$ ($i = 1, 2, \dots, 5$), which are, at this level, not plagued with chiral anomalies. The $\mathbb{Z}_4$ symmetry

$$\mathcal{D}_1 : \begin{cases} E_\mu \rightarrow iE_\mu, & E_\tau \rightarrow iE_\tau, \\ \phi_3 \rightarrow -i\phi_3, & \phi_4 \rightarrow -i\phi_4, \\ \phi_5 \rightarrow -i\phi_5, & \phi_6 \rightarrow -i\phi_6, \\ \phi_7 \rightarrow -i\phi_7, & \phi_8 \rightarrow -i\phi_8 \end{cases} \quad (1)$$

forbids dimension-four Yukawa coupling terms in the μ - τ -subsector of the charged lepton sector. The $\mathbb{Z}_2$ symmetry

$$\mathcal{D}_2 : \begin{cases} E_e \rightarrow -E_e, \\ \phi_1 \rightarrow -\phi_1, & \phi_2 \rightarrow -\phi_2 \end{cases} \quad (2)$$

sets to leading order the e - e -element of the charged lepton mass matrix equal to zero. It has been pointed out that bimaximal leptonic mixing corresponds to a permutation symmetry of the second and third generation [16]. Thus, we will introduce the three permutation symmetries

$$\mathcal{D}_3 : \begin{cases} L_\mu \rightarrow -L_\mu, & E_\mu \rightarrow -E_\mu, \\ \phi_1 \leftrightarrow \phi_2, & \phi_3 \leftrightarrow \phi_4, \end{cases} \quad (3a)$$

$$\mathcal{D}_4 : \begin{cases} L_\mu \rightarrow -L_\mu, \\ \phi_1 \leftrightarrow \phi_2, & \phi_5 \leftrightarrow \phi_6, \\ \phi_7 \rightarrow -\phi_7, \end{cases} \quad (3b)$$

$$\mathcal{D}_5 : \begin{cases} L_\mu \leftrightarrow L_\tau, & E_\mu \leftrightarrow E_\tau, \\ \phi_2 \rightarrow -\phi_2, & \phi_4 \rightarrow -\phi_4, \\ \phi_6 \rightarrow -\phi_6, & \phi_7 \leftrightarrow \phi_8. \end{cases} \quad (3c)$$

Then, the most general charged lepton mass terms, which are invariant under all symmetry transformations of our model, are given by the higher-dimensional operators

$$\mathcal{L} = \overline{L_\alpha} H_2 [(Y_{\text{eff}}^1)_{\alpha\beta} + (Y_{\text{eff}}^2)_{\alpha\beta}] E_\beta + \text{h.c.}, \quad (4)$$

where the relevant effective Yukawa interaction matrices Y_{eff}^1 and Y_{eff}^2 are on the forms

$$Y_{\text{eff}}^1 = \begin{pmatrix} 0 & B(\phi_3 - \phi_4) & B(\phi_3 + \phi_4) \\ A(\phi_1 - \phi_2) & C(\phi_5 - \phi_6) & 0 \\ A(\phi_1 + \phi_2) & 0 & C(\phi_5 + \phi_6) \end{pmatrix}, \quad (5a)$$

$$Y_{\text{eff}}^2 = \text{diag}(0, D\phi_7, D\phi_8). \quad (5b)$$

Here, the dimensionful coefficients A , B , C , and D are given by

$$A = Y_a \frac{\theta^2}{M_1^3}, \quad (6a)$$

$$B = Y_b \frac{\theta^2}{M_1^3}, \quad (6b)$$

$$C = Y_c \frac{1}{M_1}, \quad (6c)$$

$$D = Y_d \frac{\theta}{M_1^2}, \quad (6d)$$

where the quantities Y_a , Y_b , Y_c , and Y_d are arbitrary order unity coefficients and M_1 is the high mass scale of the intermediate Froggatt-Nielsen states. Actually, in Sec. 2.3, the mass scale M_1 will be related to the breakdown scale of the extra symmetries by a small expansion parameter.

2.2. The scalar potential

The most general renormalizable scalar potential, involving only the fields ϕ_i ($i = 1, 2, \dots, 6$), which is invariant under transformations of the horizontal symmetries given in Sec. 2.1, reads

$$\begin{aligned} V = & \mu_1^2 (|\phi_1|^2 + |\phi_2|^2) \\ & + \mu_2^2 (|\phi_3|^2 + |\phi_4|^2) \\ & + \mu_3^2 (|\phi_5|^2 + |\phi_6|^2) \\ & + \kappa_1^2 (|\phi_1^\dagger \phi_1|^2 + |\phi_2^\dagger \phi_2|^2) \\ & + \kappa_2^2 (|\phi_3^\dagger \phi_3|^2 + |\phi_4^\dagger \phi_4|^2) \\ & + \kappa_3^2 (|\phi_5^\dagger \phi_5|^2 + |\phi_6^\dagger \phi_6|^2) \\ & + a|\phi_1^\dagger \phi_2|^2 + b|\phi_3^\dagger \phi_4|^2 + c|\phi_5^\dagger \phi_6|^2 \\ & + d (|\phi_1^\dagger \phi_3|^2 + |\phi_2^\dagger \phi_3|^2 + |\phi_1^\dagger \phi_4|^2 + |\phi_2^\dagger \phi_4|^2) \\ & + e (|\phi_1^\dagger \phi_5|^2 + |\phi_2^\dagger \phi_5|^2 + |\phi_1^\dagger \phi_6|^2 + |\phi_2^\dagger \phi_6|^2) \\ & + f (|\phi_3^\dagger \phi_5|^2 + |\phi_4^\dagger \phi_5|^2 + |\phi_3^\dagger \phi_6|^2 + |\phi_4^\dagger \phi_6|^2) \end{aligned}$$

$$\begin{aligned}
& + \lambda_1 \left[\left(\phi_1^\dagger \phi_2 \right)^2 + \left(\phi_2^\dagger \phi_1 \right)^2 \right] \\
& + \lambda_2 \left[\left(\phi_3^\dagger \phi_4 \right)^2 + \left(\phi_4^\dagger \phi_3 \right)^2 \right] \\
& + \lambda_3 \left[\left(\phi_5^\dagger \phi_6 \right)^2 + \left(\phi_6^\dagger \phi_5 \right)^2 \right] \\
& + m_1 \left(\phi_1^\dagger \phi_2 + \phi_2^\dagger \phi_1 \right) \left(\phi_3^\dagger \phi_4 + \phi_4^\dagger \phi_3 \right) \\
& + m_2 \left(\phi_1^\dagger \phi_2 + \phi_2^\dagger \phi_1 \right) \left(\phi_5^\dagger \phi_6 + \phi_6^\dagger \phi_5 \right) \\
& + m_3 \left(\phi_3^\dagger \phi_4 + \phi_4^\dagger \phi_3 \right) \left(\phi_5^\dagger \phi_6 + \phi_6^\dagger \phi_5 \right),
\end{aligned} \tag{7}$$

where all coefficients are real. Due to the symmetries of our model, the remaining scalar fields enter relevant terms in the potential V only via quartic couplings in form of absolute squares of these fields, which means that they can be combined into the coefficients μ_i ($i = 1, 2, 3$). Therefore, we can choose the coefficients in the potential to fulfill $\mu_i^2 < 0$ ($i = 1, 2, 3$), $\kappa_i > 0$ ($i = 1, 2, 3$), and $a, b, \dots, f > 0$, which yield after spontaneous symmetry breaking (SSB) non-vanishing vacuum expectation values (VEVs) that satisfy

$$|\langle \phi_1 \rangle| = |\langle \phi_2 \rangle|, \quad |\langle \phi_3 \rangle| = |\langle \phi_4 \rangle|, \quad |\langle \phi_5 \rangle| = |\langle \phi_6 \rangle|.$$

If $\lambda_i < 0$ ($i = 1, 2, 3$), then we obtain pairwise relatively real VEVs, *i.e.*,

$$\frac{\langle \phi_1 \rangle}{\langle \phi_2 \rangle}, \frac{\langle \phi_3 \rangle}{\langle \phi_4 \rangle}, \frac{\langle \phi_5 \rangle}{\langle \phi_6 \rangle} \in \{-1, 1\}.$$

Next, choosing $m_1 < 0$ and $m_2, m_3 > 0$, we obtain

$$\frac{\langle \phi_1 \rangle}{\langle \phi_2 \rangle} \frac{\langle \phi_3 \rangle}{\langle \phi_4 \rangle} = 1 \quad \text{and} \quad \frac{\langle \phi_1 \rangle}{\langle \phi_2 \rangle} \frac{\langle \phi_5 \rangle}{\langle \phi_6 \rangle} = -1,$$

i.e., the relative sign between $\langle \phi_1 \rangle$ and $\langle \phi_2 \rangle$ is equal to the relative sign between $\langle \phi_3 \rangle$ and $\langle \phi_4 \rangle$ and opposite to the relative sign between $\langle \phi_5 \rangle$ and $\langle \phi_6 \rangle$. In Sec. 2.3, we will show that this alignment mechanism reconciles the permutation symmetry $\mathcal{D}_5$ with an approximate diagonal form of the charged lepton mass matrix.

2.3. The charged lepton mass matrix

Suppose that the SM singlet scalar fields acquire their VEVs at a high mass scale and thereby give rise to a small expansion parameter

$$\epsilon \simeq \frac{\langle \phi_i \rangle}{M_1} \simeq \frac{\langle \theta \rangle}{M_1} \simeq 10^{-1}, \tag{8}$$

where $i = 1, 2, \dots, 8$. Such small hierarchies can arise from large hierarchies in supersymmetric theories when the scalar fields acquire their VEVs along a “D-flat” direction [17]. As a consequence of the permutation symmetries $\mathcal{D}_3$, $\mathcal{D}_4$, and $\mathcal{D}_5$, the lowest energy state is two-fold degenerate. Applying the results of Sec. 2.2 and inserting Eq. (8) into Eqs. (6) and the result thereof into Eqs. (5), we obtain the two possible charged lepton mass matrices

$$M_\ell \simeq m_\tau \begin{pmatrix} 0 & \epsilon^2 & 0 \\ \epsilon^2 & \epsilon & 0 \\ 0 & 0 & 1 \end{pmatrix} \tag{9}$$

and

$$M_\ell \simeq m_\tau \begin{pmatrix} 0 & 0 & \epsilon^2 \\ 0 & 1 & 0 \\ \epsilon^2 & 0 & \epsilon \end{pmatrix}, \tag{10}$$

where m_τ is the tau mass and only the order of magnitude of the matrix elements have been indicated. Note that a permutation of the second and third generations, $L_\mu \leftrightarrow L_\tau$, $E_\mu \leftrightarrow E_\tau$, leads from one solution to the other. Let us take the first one. Diagonalization of M_ℓ [in Eq. (9)] then gives for the charged lepton masses the order-of-magnitude relations³

$$m_e/m_\mu \simeq \epsilon^2 \simeq 10^{-2} \quad \text{and} \quad m_\mu/m_\tau \simeq \epsilon \simeq 10^{-1},$$

which approximately fit the experimentally observed values [*i.e.*, $(m_e/m_\mu)_{\text{exp}} \simeq 4.8 \cdot 10^{-3}$ and $(m_\mu/m_\tau)_{\text{exp}} \simeq 5.9 \cdot 10^{-2}$ [12]]. Here m_e and m_μ are the electron and muon masses, respectively. The charged lepton mass matrix M_ℓ is diagonalized by a rotation of the left-handed charged lepton fields in the 1-2-plane by an angle $\theta_{12}^\ell \simeq 6^\circ$

³The charged lepton mass spectrum is: $m_e \equiv |\lambda_1|$, $m_\mu \equiv |\lambda_2|$, $m_\tau \equiv |\lambda_3|$, where $\lambda_1 = \frac{\epsilon}{2} (1 - \sqrt{1 + 4\epsilon^2}) m_\tau \simeq -\epsilon^3 m_\tau$, $\lambda_2 = \frac{\epsilon}{2} (1 + \sqrt{1 + 4\epsilon^2}) m_\tau \simeq \epsilon m_\tau$, and $\lambda_3 = m_\tau$ are the eigenvalues of the matrix M_ℓ .

($\theta_{12}^\ell = \epsilon - \frac{4}{3}\epsilon^3 + \mathcal{O}(\epsilon^5)$, where $\epsilon \simeq 0.1$, $\theta_{13}^\ell = 0$, and $\theta_{23}^\ell = 0$), which will finally give a contribution to all leptonic mixing angles (see App. A).

3. The neutrino mass matrix

Let us now turn our discussion to the neutrino mass matrix. As intermediate Froggatt–Nielsen states we will assume two heavy SM singlet Dirac fermions F_1 and F_2 , which have masses of the order M_1 . In order to account for the smallness of the neutrino masses, we will furthermore introduce three SM singlet Dirac fermions N_e , N_μ , and N_τ , which have masses of the order of some relevant high mass scale M_2 . From the assignment of the charges (Q_1, Q_2) to the lepton doublets, the structure of the (Q_1, Q_2) charges associated with the effective neutrino mass matrix follows immediately:

		L_e	L_μ	L_τ
	(Q_1, Q_2)	$(1, 0)$	$(0, 1)$	$(0, 1)$
$\overline{L}_e^c$	$(1, 0)$	$(2, 0)$	$(1, 1)$	$(1, 1)$
$\overline{L}_\mu^c$	$(0, 1)$	$(1, 1)$	$(0, 2)$	$(0, 2)$
$\overline{L}_\tau^c$	$(0, 1)$	$(1, 1)$	$(0, 2)$	$(0, 2)$

Note that the given representation content so far forbids any neutrino mass term. We therefore introduce the additional SM singlet scalar fields ϕ_9 , ϕ_{10} , ϕ_{11} , and ϕ_{12} and we assign the charges Q_1 , Q_2 , and Q_3 to the fields as follows:

	(Q_1, Q_2, Q_3)
N_e	$(1, 0, 0)$
N_μ, N_τ	$(0, 1, 0)$
F_1	$(1, 0, 0)$
F_2	$(-1, 0, 1)$
ϕ_9, ϕ_{10}	$(-1, -1, 0)$
ϕ_{11}	$(-2, 0, 1)$
ϕ_{12}	$(0, 0, 0)$

Note that our model is again kept free from chiral anomalies, since the heavy fermions are vector representations under transformations of all U(1) charges. The Dirac neutrino fields and the scalar fields transform under the discrete symmetries $\mathcal{D}_3$, $\mathcal{D}_4$, $\mathcal{D}_5$, and the additional discrete sym-




Figure 1. The dimension six operator for $\alpha = \mu, \tau$ and $\phi_\mu \equiv \phi_9$, $\phi_\tau \equiv \phi_{10}$, generating the e - μ - and e - τ -elements in the effective neutrino mass matrix.




Figure 2. The dimension eight operators, generating the e - e -element in the effective neutrino mass matrix.

metry $\mathcal{D}_6$ as

$$\mathcal{D}_3 : \dots, N_\mu \rightarrow -N_\mu, \phi_9 \rightarrow -\phi_9, \quad (11a)$$

$$\mathcal{D}_4 : \dots, N_\mu \rightarrow -N_\mu, \phi_9 \rightarrow -\phi_9, \quad (11b)$$

$$\mathcal{D}_5 : \dots, N_\mu \leftrightarrow N_\tau, \phi_9 \leftrightarrow \phi_{10}, \quad (11c)$$

$$\mathcal{D}_6 : \begin{cases} N_e \rightarrow iN_e, \\ \phi_{11} \rightarrow -i\phi_{11}, \quad \phi_{12} \rightarrow i\phi_{12}. \end{cases} \quad (11d)$$

It is easily verified that the results for the charged lepton mass matrix remain unchanged by this new representation content. The leading order tree-level realizations of the higher-dimensional operators, which generate the neutrino masses, are shown in Figs. 1 and 2. After SSB, the effective neutrino mass matrix M_ν will be on the approximate bimaximal mixing form [11]

$$M_\nu = \begin{pmatrix} A' & B' & -B' \\ B' & 0 & 0 \\ -B' & 0 & 0 \end{pmatrix}, \quad (12)$$

where

$$A' = Y'_a \frac{\langle H_1 \rangle^2}{M_2} \frac{\langle \phi_{11} \rangle \langle \phi_{12} \rangle \langle \theta \rangle}{(M_1)^3}, \quad (13a)$$

$$B' = Y'_b \frac{\langle H_1 \rangle^2}{M_2} \frac{\langle \phi_9 \rangle}{M_1}, \quad (13b)$$

and the quantities Y'_a and Y'_b are arbitrary order unity coefficients. Note that the permutation symmetry $\mathcal{D}_5$ establishes $|\langle \phi_9 \rangle| = |\langle \phi_{10} \rangle|$, and hence, the magnitudes of the e - μ - and e - τ -entries in the effective neutrino mass matrix M_ν are exactly degenerate. The possible relative phase φ between these entries can be eliminated by, *e.g.*, the field redefinition $L_\tau \rightarrow e^{i\varphi} L_\tau$, since it does not affect the mixing angles in the charged lepton sector. If we assume that all SM singlet scalar fields acquire their VEVs at the breakdown scale of the additional horizontal symmetries, then Eq. (8) also applies to the indices $i = 9, 10, 11, 12$. Therefore, we can parameterize the effective neutrino mass matrix M_ν with the same expansion parameter ϵ that was used for the charged lepton mass matrix M_ℓ and we find the relative suppression ratio $A'/B' \simeq \epsilon^2 \simeq 10^{-2}$ from Eq. (13). Thus, diagonalizing the neutrino mass matrix M_ν in Eq. (12), we obtain the following spectrum of the neutrino masses⁴

$$m_1 \simeq m_2 \quad \text{and} \quad m_3 = 0,$$

which is on the inverted hierarchical form (*i.e.*, $m_3 \ll m_1 \simeq m_2 \Rightarrow 0 \simeq |\Delta m_{21}^2| \ll |\Delta m_{32}^2| \simeq |\Delta m_{31}^2|$). Using the MSW LMA and atmospheric neutrino mass squared differences⁵ [9,3]

$$\begin{aligned} \Delta m_\odot^2 &\equiv |\Delta m_{21}^2| \simeq 3.7 \cdot 10^{-5} \text{ eV}^2 \sim 10^{-5} \text{ eV}^2, \\ \Delta m_{\text{atm}}^2 &\equiv |\Delta m_{32}^2| \simeq 2.5 \cdot 10^{-3} \text{ eV}^2 \sim 10^{-3} \text{ eV}^2, \end{aligned}$$

and a VEV of the order of the electroweak scale $\langle H_1 \rangle \simeq 10^2 \text{ GeV}$, we obtain from Eqs. (13) the high mass scale $M_2 \simeq 10^{14} \text{ GeV}$ (as well as $m_1 \simeq m_2 \simeq 0.05 \text{ eV}$ and $\epsilon \simeq 0.1$). The entry A' of the matrix in Eq. (12) induces a deviation from

⁴The neutrino mass spectrum is: $m_1 \equiv |\lambda'_1|$, $m_2 \equiv |\lambda'_2|$, $m_3 \equiv |\lambda'_3|$, where $\lambda'_1 \simeq -B'\sqrt{2}$, $\lambda'_2 \simeq B'\sqrt{2}$, and $\lambda'_3 = 0$ are the eigenvalues of the matrix M_ν .

⁵Here $\Delta m_{ab}^2 = \lambda'^2_a - \lambda'^2_b$, *i.e.*, $|\Delta m_{21}^2| \simeq 2\sqrt{2}|A'B'|$ and $|\Delta m_{32}^2| \simeq |\Delta m_{31}^2| \simeq 2|B'|^2$.

maximal solar mixing, which is of the order 0.1° ($\theta_{12}^\nu = \frac{\pi}{4} + \frac{1}{4\sqrt{2}}\epsilon^2 - \frac{1}{96\sqrt{2}}\epsilon^4 + \mathcal{O}(\epsilon^{10})$, where $\epsilon \simeq \sqrt{A'/B'} \simeq 0.1$, $\theta_{13}^\nu = 0$, and $\theta_{23}^\nu = 45^\circ$). This deviation can be neglected in comparison with the contribution coming from the charged lepton sector, which is about 6° (see Sec. 2.3), resulting in a change of the mixing angles θ_{12} and θ_{13} by approximately 4° , while the atmospheric mixing angle θ_{23} practically stays maximal (see App. A). Thus, the leptonic mixing angles are

$$\theta_{12} \simeq 41^\circ, \quad \theta_{13} \simeq 4^\circ, \quad \text{and} \quad \theta_{23} \simeq 45^\circ.$$

Hence, our model predicts the charged lepton mass hierarchy spectrum, an inverted neutrino mass hierarchy spectrum, bilarge leptonic mixing, as well as it reproduces the mass squared differences to lie within the ranges preferred by the MSW LMA solution⁶ and atmospheric neutrino data⁷. In particular, it yields a significant deviation from maximal solar mixing. However, the solar mixing angle is bounded from below by approximately 41° and it is therefore still too close to maximal to be in the 95% (or 99.73%) confidence level region of the MSW LMA solution⁸ [8,9].

4. Summary and conclusions

In summary, we have presented a model built upon extra fields, horizontal (flavor) symmetries, and higher-dimensional operators including the Froggatt–Nielsen mechanism. This model naturally yields the well-known mass matrix textures

$$\begin{pmatrix} 0 & \epsilon^2 & 0 \\ \epsilon^2 & \epsilon & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \epsilon^2 & 1 & -1 \\ 1 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}$$

for the charged leptons and the neutrinos, respectively, which involve the same small expansion parameter $\epsilon \simeq 0.1$. These textures reproduce the mass hierarchy among the charged leptons very

⁶@ 99.73% C.L.: $1.9 \cdot 10^{-5} \text{ eV}^2 \lesssim \Delta m_\odot^2 \lesssim 2.7 \cdot 10^{-4} \text{ eV}^2$ [18]; best-fit: $\Delta m_\odot^2 \simeq 3.7 \cdot 10^{-5} \text{ eV}^2$ [9]

⁷@ 90% C.L.: $1.6 \cdot 10^{-3} \text{ eV}^2 \lesssim \Delta m_{\text{atm}}^2 \lesssim 4.0 \cdot 10^{-3} \text{ eV}^2$; best-fit: $\Delta m_{\text{atm}}^2 \simeq 2.5 \cdot 10^{-3} \text{ eV}^2$ [3]

⁸@ 99.73% C.L.: $0.22 \lesssim \tan^2 \theta_{12} \lesssim 0.71 \Rightarrow 25^\circ \lesssim |\theta_{12}| \lesssim 40^\circ$ [18]; best-fit: $\tan^2 \theta_{12} \simeq 0.37 \Rightarrow |\theta_{12}| \simeq 31^\circ$ [9]

accurately as well as they give rise to the most probable values of the mass squared differences for the neutrinos coming from solar and atmospheric neutrino data. In addition, assuming no $\mathcal{CP}$ violation, *i.e.*, $\delta = 0$, the model gives bilarge leptonic mixing, *i.e.*, $\theta_{12} \simeq 41^\circ$, $\theta_{13} \simeq 4^\circ$, and $\theta_{23} \simeq 45^\circ$, which is in very good agreement with the present experimental data. The mixing angle θ_{12} is on the borderline of being compatible with the MSW LMA solution, which has, however, been further strengthened by recent SNO results [10] and which will, nevertheless, soon be even better tested by the KamLAND experiment [19], whereas θ_{13} is well below the CHOOZ upper bound, and θ_{23} fits perfectly the best-fit value from the Super-Kamiokande collaboration of their atmospheric neutrino data.

As a final conclusion, we have shown in App. A that the leptonic mixing angles are all dependent on the “solar” mixing angle θ_{12}^ℓ in the charged lepton sector in a non-trivial way.

Acknowledgments

We would like to thank Martin Freund, Patrick Huber, and Manfred Lindner for useful discussions.

This work was supported by the Swedish Foundation for International Cooperation in Research and Higher Education (STINT) [T.O.], the Wenner-Gren Foundations [T.O.], the Magnus Bergvall Foundation (Magn. Bergvalls Stiftelse) [T.O.], and the “Sonderforschungsbereich 375 für Astro-Teilchenphysik der Deutschen Forschungsgemeinschaft” [T.O. and G.S.].

A. The leptonic mixing matrix - A special case

The leptonic mixing matrix can be written as

$$\begin{aligned} U &= O_{23}(\theta_{23})U_{13}(\theta_{13}, \delta)O_{12}(\theta_{12}) \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & C_{23} & S_{23} \\ 0 & -S_{23} & C_{23} \end{pmatrix} \\ &\times \begin{pmatrix} C_{13} & 0 & S_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -S_{13}e^{i\delta} & 0 & C_{13} \end{pmatrix} \\ &\times \begin{pmatrix} C_{12} & S_{12} & 0 \\ -S_{12} & C_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \end{aligned} \quad (14)$$

where $C_{ab} \equiv \cos \theta_{ab}$, $S_{ab} \equiv \sin \theta_{ab}$, and $O_{ab}(\theta_{ab})$ is a rotation by an angle θ_{ab} in the ab -plane. If $\delta = 0$, then $U_{13}(\theta_{13}, 0) = O_{13}(\theta_{13})$. Assuming that all the $\mathcal{CP}$ phases are zero, *i.e.*, $\delta = \delta^\ell = \delta^\nu = 0$, then the charged lepton and neutrino mixing matrices, U^ℓ and U^ν , can, of course, also be written in the same form as the above complete leptonic mixing matrix, U , *i.e.*, we have

$$U^\ell = O_{23}^\ell(\theta_{23}^\ell) \underbrace{U_{13}^\ell(\theta_{13}^\ell, \delta^\ell)}_{O_{13}^\ell(\theta_{13}^\ell)} O_{12}^\ell(\theta_{12}^\ell), \quad (15a)$$

$$U^\nu = O_{23}^\nu(\theta_{23}^\nu) \underbrace{U_{13}^\nu(\theta_{13}^\nu, \delta^\nu)}_{O_{13}^\nu(\theta_{13}^\nu)} O_{12}^\nu(\theta_{12}^\nu). \quad (15b)$$

Thus, inserting Eqs. (15) into the definition of the leptonic mixing matrix, $U = (U_{\alpha a} \equiv U^{\ell\dagger} U^\nu)$, we find that

$$\begin{aligned} U &= O_{12}^\ell(\theta_{12}^\ell)^T O_{13}^\ell(\theta_{13}^\ell)^T O_{23}^\ell(\theta_{23}^\ell)^T \\ &\times O_{23}^\nu(\theta_{23}^\nu) O_{13}^\nu(\theta_{13}^\nu) O_{12}^\nu(\theta_{12}^\nu). \end{aligned} \quad (16)$$

Furthermore, assuming that we have only a small mixing coming from the mixing angle θ_{12}^ℓ in the charged lepton sector ($\theta_{13}^\ell = 0$, $\theta_{23}^\ell = 0$) and bi-maximal mixing in the neutrino sector ($\theta_{12}^\nu = 45^\circ$, $\theta_{13}^\nu = 0$, $\theta_{23}^\nu = 45^\circ$), we then obtain

$$\begin{aligned} U &= O_{12}^\ell(\theta_{12}^\ell)^T O_{23}^\nu(\theta_{23}^\nu = 45^\circ) O_{12}^\nu(\theta_{12}^\nu = 45^\circ) \\ &= \begin{pmatrix} C_{12}^\ell/\sqrt{2} + S_{12}^\ell/2 & C_{12}^\ell/\sqrt{2} - S_{12}^\ell/2 & -S_{12}^\ell/\sqrt{2} \\ S_{12}^\ell/\sqrt{2} - C_{12}^\ell/2 & S_{12}^\ell/\sqrt{2} + C_{12}^\ell/2 & C_{12}^\ell/\sqrt{2} \\ 1/2 & -1/2 & 1/\sqrt{2} \end{pmatrix}. \end{aligned} \quad (17)$$

The mixing angles (in the standard parameterization) of a 3×3 orthogonal mixing matrix can be read off as follows [20]:

$$\theta_{12} = \arctan \frac{U_{e2}}{U_{e1}}, \quad (18)$$

$$\theta_{13} = \arcsin U_{e3}, \quad (19)$$

$$\theta_{23} = \arctan \frac{U_{\mu 3}}{U_{\tau 3}}. \quad (20)$$

Thus, inserting the appropriate matrix elements of the matrix U in Eq. (17) into Eqs. (18) - (20), we finally obtain

$$\theta_{12} = \arctan \frac{\cos \theta_{12}^\ell - \frac{1}{\sqrt{2}} \sin \theta_{12}^\ell}{\cos \theta_{12}^\ell + \frac{1}{\sqrt{2}} \sin \theta_{12}^\ell}, \quad (21)$$

$$\theta_{13} = -\arcsin \left(\frac{1}{\sqrt{2}} \sin \theta_{12}^\ell \right), \quad (22)$$

$$\theta_{23} = \arctan \cos \theta_{12}^\ell. \quad (23)$$

When θ_{12}^ℓ is small ($\theta_{12}^\ell \ll 1$), we have⁹

$$\theta_{12} = \frac{\pi}{4} - \frac{1}{\sqrt{2}} \theta_{12}^\ell - \frac{1}{6\sqrt{2}} \theta_{12}^{\ell 3} + \mathcal{O}(\theta_{12}^{\ell 5}), \quad (24)$$

$$\theta_{13} = -\frac{1}{\sqrt{2}} \theta_{12}^\ell + \frac{1}{12\sqrt{2}} \theta_{12}^{\ell 3} + \mathcal{O}(\theta_{12}^{\ell 5}), \quad (25)$$

$$\theta_{23} = \frac{\pi}{4} - \frac{1}{4} \theta_{12}^{\ell 2} - \frac{1}{24} \theta_{12}^{\ell 4} + \mathcal{O}(\theta_{12}^{\ell 6}). \quad (26)$$

Note that all leptonic mixing angles receive contribution from the small mixing angle θ_{12}^ℓ in the charged lepton sector. Furthermore, we observe that θ_{12} has first order corrections in θ_{12}^ℓ , whereas θ_{23} has only second order corrections in θ_{12}^ℓ . The mixing angle θ_{13} is directly proportional to the mixing angle θ_{12}^ℓ (when θ_{12}^ℓ is small), which means that if θ_{12}^ℓ is small, then θ_{13} will also be small. In fact, $|\theta_{12}^\ell| \lesssim 13.1^\circ$ has to be fulfilled in order for the mixing angle θ_{13} to be below the CHOOZ upper bound $\sin^2 2\theta_{13} \lesssim 0.10$ (*i.e.*, $|\theta_{13}| \lesssim 9.2^\circ$) [21].

⁹Introducing a small deviation η from maximal solar mixing in the neutrino sector (*i.e.*, $\theta_{12}^\nu = 45^\circ \rightarrow \theta_{12}^\nu = 45^\circ + \eta$, where $\eta \ll 1$), we find that $\theta_{12} = \frac{\pi}{4} + \eta - \frac{1}{\sqrt{2}} \theta_{12}^\ell - \frac{1}{6\sqrt{2}} \theta_{12}^{\ell 3} + \mathcal{O}(\theta_{12}^{\ell 5})$, whereas θ_{13} and θ_{23} remain unchanged.

REFERENCES

1. E. Witten, Nucl. Phys. B (Proc. Suppl.) 91 (2000) 3, [hep-ph/0006332](#).
2. Super-Kamiokande Collaboration, Y. Fukuda et al., Phys. Rev. Lett. 81 (1998) 1562, [hep-ex/9807003](#); *ibid.* 82 (1999) 2644, [hep-ex/9812014](#); S. Fukuda et al., Phys. Rev. Lett. 85 (2000) 3999, [hep-ex/0009001](#).
3. Super-Kamiokande Collaboration, T. Toshito, [hep-ex/0105023](#).
4. Super-Kamiokande Collaboration, S. Fukuda et al., Phys. Rev. Lett. 86 (2001) 5651, [hep-ex/0103032](#); M.B. Smy, [hep-ex/0106064](#); SNO Collaboration, Q.R. Ahmad et al., Phys. Rev. Lett. 87 (2001) 071301, [nucl-ex/0106015](#); [nucl-ex/0204008](#).
5. Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. 28 (1962) 870.
6. S.P. Mikheyev and A.Y. Smirnov, Yad. Fiz. 42 (1985) 1441, [Sov. J. Nucl. Phys. 42 (1985) 913]; Nuovo Cimento C 9 (1986) 17; L. Wolfenstein, Phys. Rev. D 17 (1978) 2369.
7. G.L. Fogli et al., Phys. Rev. D 64 (2001) 093007, [hep-ph/0106247](#); A. Bandyopadhyay et al., Phys. Lett. B 519 (2001) 83, [hep-ph/0106264](#); P.I. Krastev and A.Y. Smirnov, Phys. Rev. D 65 (2002) 073022, [hep-ph/0108177](#); P. Aliani et al., [hep-ph/0111418](#).
8. J.N. Bahcall, M.C. Gonzalez-Garcia and C. Peña-Garay, J. High Energy Phys. 08 (2001) 014, [hep-ph/0106258](#).
9. J.N. Bahcall, M.C. Gonzalez-Garcia and C. Peña-Garay, J. High Energy Phys. 04 (2002) 014, [hep-ph/0111150](#).
10. SNO Collaboration, Q.R. Ahmad et al., [nucl-ex/0204009](#).
11. V. Barger et al., Phys. Lett. B 437 (1998) 107, [hep-ph/9806387](#).
12. Particle Data Group, D.E. Groom et al., Eur. Phys. J. C 15 (2000) 1, <http://pdg.lbl.gov/>.
13. S. Weinberg, Phys. Rev. Lett. 43 (1979) 1566; F. Wilczek and A. Zee, Phys. Rev. Lett. 43 (1979) 1571.
14. C.D. Froggatt and H.B. Nielsen, Nucl.

- Phys. B 147 (1979) 277.
15. K.S. Babu and C.N. Leung, Nucl. Phys. B 619 (2001) 667, [hep-ph/0106054](#); A. Pérez-Lorenzana and C.A. de S. Pires, Phys. Lett. B 522 (2001) 297, [hep-ph/0108158](#); J.C. Montero, C.A. de S. Pires and V. Pleitez, [hep-ph/0112203](#); C.D. Froggatt, H.B. Nielsen and Y. Takanishi, Nucl. Phys. B (to be published), [hep-ph/0201152](#).
 16. R.N. Mohapatra, A. Pérez-Lorenzana and C.A. de S. Pires, Phys. Lett. B 474 (2000) 355, [hep-ph/9911395](#); W. Grimus and L. Lavoura, J. High Energy Phys. 07 (2001) 045, [hep-ph/0105212](#); R.N. Mohapatra, Phys. Rev. D 64 (2001) 091301, [hep-ph/0107264](#).
 17. E. Witten, Phys. Lett. B 105 (1981) 267; M. Leurer, Y. Nir and N. Seiberg, Nucl. Phys. B 420 (1994) 468, [hep-ph/9310320](#).
 18. M.C. Gonzalez-Garcia and Y. Nir, [hep-ph/0202058](#).
 19. KamLAND Collaboration, A. Piepke, Nucl. Phys. B (Proc. Suppl.) 91 (2001) 99.
 20. T. Ohlsson and H. Snellman, J. Math. Phys. 41 (2000) 2768, [hep-ph/9910546](#), 42 (2001) 2345(E); T. Ohlsson, Phys. Scripta T93 (2001) 18.
 21. CHOOZ Collaboration, M. Apollonio et al., Phys. Lett. B 420 (1998) 397, [hep-ex/9711002](#); *ibid.* 466 (1999) 415, [hep-ex/9907037](#); C. Bemporad, Nucl. Phys. B (Proc. Suppl.) 77 (1999) 159.